Ultraviolet And Fluorescence Spectroscopy

from chapter _____ in the recommended text

A. Introduction

B. Fundamental Physics

more
X-ray

ground state energy level to a(n) *excited*

inversely proportional to the *energies*
directly related to their *energies*.

broad
X-ray
Chromophores
cross-section and the *more*

C. Molecular Orbital Diagrams Of Alkenes, Dienes, and Polyenes

more than one single


non-conj'd


conj'd


conj'd


conj'd


non-conj'd


non-conj'd

$H_2C{=}CH_2$
non-conj'd


conj'd



conj'd

conj'd

non-conj'd


conj'd


non-conj'd


conj'd


β-carotene *conj'd*


isoprene
conj'd


limonene
non-conj'd



squalene non-conj'd



arachidinoic acid non-conj'd



porphyrin conj'd

is



phosphoiodyns A
non-conj'd



debilisone
conj'd

an alternative to
n

1
bonding π- and *antibonding*
Maximal



ultraviolet region resulting in an *excited*
IR energy

larger cross-sections, therefore they absorb *more*
absorbance of the *chromophore*
larger



decreases
lower energy quanta of *increased*

vibrational energy states, while *electronic*
IR
UV quanta.

UV
IR.

IR
UV
greater than for transitions like **G**.



multiple
smaller
UV and transitions between *electronic*
IR emissions.
nano-second
fluorescent radiation
rigid molecules



sensitive
higher
higher
fluorescence spectroscopy
fluors.
less

rigid



a BODIPY



a fluorescein



a rhodamine



a coumarin

Of the probes shown above, the ones which has fluorescence that is most sensitive to reduced pH is the *fluorescein*, whereas the *BODIPY* is least sensitive to pH.

This is because *fluorescein* contain phenoate-*O*$^-$ groups that can be protonated as the pH is reduced from 7.0.

Fluorescence of the *BODIPY / fluorescein / rhodamine / coumarin* is likely to be most sensitive to the dipole moment of the solvent it is in because the oscillation of charge in this molecule is unsymmetrical.

Which of the fluor solutions below emit the highest energy light *B*, and which of them emit at the longest wavelength *V*.



Circle the correct definition of fluorescence quantum yield from the following choices:

$\frac{\text{\# photons absorbed}}{\text{\# photons emitted}}$ $\frac{\text{\# photons emitted}}{\text{\# photons absorbed}}$ $\frac{\text{\# photons lost as heat}}{\text{\# photons absorbed}}$

Circle the correct descriptor of fluor brightness:

quantum yield x absorbance at excitation wavelengths

quantum yield x absorbance at λ_{max}

absorbance at excitation wavelengths only

quantum yield only

absorbance at λ_{max} only